

Graduate Texts in Mathematics

GTM

Steven G. Krantz

Geometric Analysis of the Bergman Kernel and Metric

 Springer

Graduate Texts in Mathematics

Series Editors:

Sheldon Axler
San Francisco State University, San Francisco, CA, USA

Kenneth Ribet
University of California, Berkeley, CA, USA

Advisory Board:

Colin Adams, *Williams College, Williamstown, MA, USA*

Alejandro Adem, *University of British Columbia, Vancouver, BC, Canada*

Ruth Charney, *Brandeis University, Waltham, MA, USA*

Irene M. Gamba, *The University of Texas at Austin, Austin, TX, USA*

Roger E. Howe, *Yale University, New Haven, CT, USA*

David Jerison, *Massachusetts Institute of Technology, Cambridge, MA, USA*

Jeffrey C. Lagarias, *University of Michigan, Ann Arbor, MI, USA*

Jill Pipher, *Brown University, Providence, RI, USA*

Fadil Santosa, *University of Minnesota, Minneapolis, MN, USA*

Amie Wilkinson, *University of Chicago, Chicago, IL, USA*

Graduate Texts in Mathematics bridge the gap between passive study and creative understanding, offering graduate-level introductions to advanced topics in mathematics. The volumes are carefully written as teaching aids and highlight characteristic features of the theory. Although these books are frequently used as textbooks in graduate courses, they are also suitable for individual study.

For further volumes:

<http://www.springer.com/series/136>

Steven G. Krantz

Geometric Analysis of the Bergman Kernel and Metric

 Springer

Steven G. Krantz
Department of Mathematics
Washington University at St. Louis
St. Louis, MO, USA

ISSN 0072-5285
ISBN 978-1-4614-7923-9 ISBN 978-1-4614-7924-6 (eBook)
DOI 10.1007/978-1-4614-7924-6
Springer New York Heidelberg Dordrecht London

Library of Congress Control Number: 2013944372

Mathematics Subject Classification (2010): 30C40, 32A25, 32A26, 32A36, 32H40, 46C05

© Springer Science+Business Media New York 2013

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

*Dedicated to the memory of Stefan Bergman,
an extraordinarily profound and original
mathematician*

Preface

The Bergman kernel and metric have been a seminal part of geometric analysis and partial differential equations since their invention by Stefan Bergman in 1922. Applications to holomorphic mappings, to function theory, to partial differential equations, and to differential geometry have kept the techniques plugged into the mainstream of mathematics for 90 years.

The Bergman kernel is based on a very simple idea: that the square-integrable holomorphic functions on a bounded domain in that complex space form a Hilbert space. Moreover, a simple formal argument shows that Hilbert space possesses a so-called reproducing kernel. This is an integration kernel which reproduces each element of the space. The kernel has wonderful invariance properties, leading to the Bergman metric. The Bergman kernel and metric have developed into powerful tools for function theory, analysis, differential geometry, and partial differential equations. The purpose of this book is to exposit this theory (particularly in the context of several complex variables), examine its key features, and bring the reader up to speed with some of the latest developments.

Bergman wrote several books about his kernel and contributed mightily to its development. The idea caught on widely, and the kernel became a standard device in the field. The Bergman metric was the first-ever Kähler metric, and that in turn spawned the vital subject of complex differential geometry. Ahlfors (in one variable), Chern (in several variables), Greene–Wu, and many others played a decisive role in this development.

Bergman's ideas received a major boost in the 1970s when Charles Fefferman did his Fields Medal-winning work on the boundary behavior of biholomorphic mappings. The key device in his analysis was the Bergman kernel and metric. Since then, a myriad of workers, from Bell and Ligocka to Webster to Greene–Krantz, Krantz–Li, Kim–Krantz, Isaev–Krantz, and many others, have developed and extended Bergman's theory. It is now part of the lingua franca of complex analysis, and the technique of reproducing kernels which it has spawned is part of every analyst's toolkit.

Fefferman's work inspired many others to examine the utility of the Bergman theory in the study of biholomorphic mappings. Bell's condition R , formulated

in terms of regularity properties of the Bergman projection, has proved to be an influential and powerful weapon in the subject. In turn, condition R can be formulated in terms of the regularity theory of subelliptic partial differential equations, and this connection has had a key influence on the directions of research.

The Bergman metric was the first “universal” (in the sense that it can be constructed on virtually any domain) example of a metric that is invariant under biholomorphic mappings. [The Poincaré metric was of course the primordial example on the disc.] Today there are the Kobayashi–Royden metric, the Sibony metric, the Carathéodory metric, and many others. This is a useful tool in geometric analysis and function theory.

The connections of the Bergman kernel with partial differential equations, especially the extremal properties of the kernel and metric, are profound. Bergman himself explored applications of his theory to elliptic partial differential equations. Today we see the Bergman kernel as inextricably linked with the $\bar{\partial}$ -Neumann problem. This link played a vital role in Fefferman’s work and later proved crucial to Greene–Krantz and many of the other workers in the subject. Certainly Donald Spencer and J. J. Kohn were the pioneers of this symbiosis. Today the interaction is prospering. Another development is that the Bergman theory enjoys connections with the Monge–Ampère equation. That nonlinear partial differential equation contains important information about biholomorphic mappings and about the construction of geometries.

Connections with harmonic analysis are another exciting, and relatively new, aspect of the Bergman paradigm. Coifman–Rochberg–Weiss used the Bergman kernel in their proof of the H^1/BMO duality theorem on the ball, and Krantz–Li exploited it further in their study on strictly pseudoconvex domains. Many of the natural artifacts of harmonic analysis—including approach regions for Fatou theorems—are most propitiously formulated in terms of Bergman geometry or the boundary asymptotics of the kernel. The Bergman kernel is now a standard artifact of the harmonic analysis of several complex variables.

This text will in fact be a thoroughgoing treatment of all the basic analytic and geometric aspects of Bergman’s theory. This will include

- Definitions of the Bergman kernel
- Definition and basic properties of the Bergman metric
- Calculation of the Bergman kernel and metric
- Invariance properties of the kernel and metric
- Boundary asymptotics of the kernel and metric
- Asymptotic expansions for the Bergman kernel
- Applications to function theory
- Applications to geometry
- Applications to partial differential equations
- Interpretations in terms of functional analysis
- The geometry of the Bergman metric
- Curvature of the Bergman metric
- The Bergman kernel and metric on manifolds

There are a few recent treatises on the Bergman kernel, notably those by Hedenmalm–Korenblum–Zhu and Duren–Schuster. But these books concentrate on the one-variable theory; they are also oriented towards the functional analysis aspects of the Bergman kernel. Our focus instead is the geometry of several complex variables and the contexts of real analysis, complex analysis, harmonic analysis, and differential geometry. This puts Bergman’s ideas into a much broader arena and provides many more opportunities for applications and illustrations. We will certainly touch on the functional analysis properties of the Bergman projection, but these will not be our main focus. We shall also cover selected topics of the one complex variable theory. There is little overlap between this book and the two books cited above.

We would also be remiss not to mention the book of Ma and Marinescu on the Kähler geometry aspects of the Bergman theory. Certainly the Bergman metric was the very first Kähler metric, and this in turn has spawned the active and fruitful area of multivariable complex differential geometry. Various parts of the present book touch on this Kähler theory.

Lurking in the background behind Fefferman’s biholomorphic mapping theorem were Bergman representative coordinates—yet another outgrowth of the Bergman kernel and metric. This is a much-underappreciated aspect of the theory and one that we shall treat in detail in the text. In fact there are many aspects of the Bergman theory that tend only to be known to experts and are not readily accessible in the literature. We intend to treat many of those. Several of the topics in this text appear here for the first time in book form.

We intend this to be a book for students as well as seasoned researchers. All needed background will be provided. The reader is only assumed to have had a solid course in complex variables and some basic background in real and functional analysis. A little exposure to geometry will be helpful, but is not a requirement. There are many illustrative examples and some useful figures. The book abounds with useful and instructive calculations, many of which cannot be found elsewhere. Every chapter ends with a selection of exercises, which should serve to help the reader get more directly involved in the subject matter. It will cause him/her to consult the literature, to calculate, and to learn by doing.

The book will help the novice reader to see how analysis is used in practice and how it can be evolved into a seminal tool for research. It is important for the student to see fundamental mathematics used *in vitro* in order to understand how research develops and grows.

It is a pleasure to thank E. M. Stein for introducing me to the Bergman kernel and Robert E. Greene for teaching me the geometric aspects. I have had many collaborators in my study of the kernel, and I offer them all my gratitude. I thank my editor, Elizabeth Loew, for her constant enthusiasm and support. And I thank the several referees for this book, who contributed a wealth of ideas and information.

Contents

| | | |
|----------|--|----|
| 1 | Introductory Ideas | 1 |
| 1.1 | The Bergman Kernel | 1 |
| 1.1.1 | Calculating the Bergman Kernel | 14 |
| 1.1.2 | The Poincaré-Bergman Distance on the Disc | 19 |
| 1.1.3 | Construction of the Bergman Kernel by Way of Differential Equations | 20 |
| 1.1.4 | Construction of the Bergman Kernel by Way of Conformal Invariance | 23 |
| 1.2 | The Szegő and Poisson–Szegő Kernels | 25 |
| 1.3 | Formal Ideas of Aronszajn | 35 |
| 1.4 | A New Bergman Basis | 36 |
| 1.5 | Further Examples | 39 |
| 1.6 | A Real Bergman Space | 40 |
| 1.7 | The Behavior of the Singularity in a General Setting | 41 |
| 1.8 | The Annulus | 43 |
| 1.9 | A Direct Connection Between the Bergman and Szegő Kernels | 45 |
| 1.9.1 | Introduction | 45 |
| 1.9.2 | The Case of the Disc | 45 |
| 1.9.3 | The Unit Ball in \mathbb{C}^n | 49 |
| 1.9.4 | Strongly Pseudoconvex Domains | 52 |
| 1.9.5 | Concluding Remarks | 53 |
| 1.10 | Multiply Connected Domains | 53 |
| 1.11 | The Bergman Kernel for a Sobolev Space | 54 |
| 1.12 | Ramadanov’s Theorem | 56 |
| 1.13 | Coda on the Szegő Kernel | 58 |
| 1.14 | Boundary Localization | 59 |
| 1.14.1 | Definitions and Notation | 60 |
| 1.14.2 | A Representative Result | 60 |

| | | |
|----------|--|-----|
| 1.14.3 | The More General Result in the Plane | 62 |
| 1.14.4 | Domains in Higher-Dimensional Complex Space..... | 62 |
| | Exercises | 65 |
| 2 | The Bergman Metric | 71 |
| 2.1 | Smoothness to the Boundary of Biholomorphic Mappings | 71 |
| 2.2 | Boundary Behavior of the Bergman Metric | 81 |
| 2.3 | The Biholomorphic Inequivalence of the Ball and the Polydisc | 83 |
| | Exercises | 84 |
| 3 | Further Geometric and Analytic Theory | 87 |
| 3.1 | Bergman Representative Coordinates..... | 87 |
| 3.2 | The Berezin Transform | 90 |
| 3.2.1 | Preliminary Remarks | 90 |
| 3.2.2 | Introduction to the Poisson–Bergman Kernel | 91 |
| 3.2.3 | Boundary Behavior | 94 |
| 3.3 | Ideas of Fefferman..... | 98 |
| 3.4 | Results on the Invariant Laplacian | 100 |
| 3.5 | The Dirichlet Problem for the Invariant Laplacian on the Ball | 109 |
| 3.6 | Concluding Remarks | 115 |
| | Exercises | 115 |
| 4 | Partial Differential Equations | 117 |
| 4.1 | The Idea of Spherical Harmonics | 117 |
| 4.2 | Advanced Topics in the Theory of Spherical Harmonics: The Zonal Harmonics | 117 |
| 4.3 | Spherical Harmonics in the Complex Domain and Applications.... | 130 |
| 4.4 | An Application to the Bergman Projection | 141 |
| | Exercises | 145 |
| 5 | Further Geometric Explorations | 147 |
| 5.1 | Introductory Remarks | 147 |
| 5.2 | Semicontinuity of Automorphism Groups | 151 |
| 5.3 | Convergence of Holomorphic Mappings | 156 |
| 5.3.1 | Finite Type in Dimension Two | 156 |
| 5.4 | The Semicontinuity Theorem | 166 |
| 5.5 | Some Examples | 168 |
| 5.6 | Further Remarks | 168 |
| 5.7 | The Lu Qi-Keng Conjecture | 169 |
| 5.8 | The Lu Qi-Keng Theorem..... | 171 |
| 5.9 | The Dimension of the Bergman Space | 174 |
| 5.10 | The Bergman Theory on a Manifold..... | 178 |
| 5.10.1 | Kernel Forms | 178 |
| 5.10.2 | The Invariant Metric | 182 |
| 5.11 | Boundary Behavior of the Bergman Metric | 184 |
| | Exercises | 185 |

| | |
|--|-----|
| 6 Additional Analytic Topics | 187 |
| 6.1 The Diederich–Fornæss Worm Domain | 187 |
| 6.2 More on the Worm | 192 |
| 6.3 Non-Smooth Versions of the Worm Domain | 199 |
| 6.4 Irregularity of the Bergman Projection | 200 |
| 6.5 Irregularity Properties of the Bergman Kernel | 205 |
| 6.6 The Kohn Projection Formula | 207 |
| 6.7 Boundary Behavior of the Bergman Kernel | 208 |
| 6.7.1 Hörmander’s Result on Boundary Behavior | 209 |
| 6.7.2 The Fefferman’s Asymptotic Expansion | 215 |
| 6.8 The Bergman Kernel for a Sobolev Space | 221 |
| 6.9 Regularity of the Dirichlet Problem on a Smoothly Bounded Domain and Conformal Mapping | 224 |
| 6.10 Existence of Certain Smooth Plurisubharmonic Defining Functions for Strictly Pseudoconvex Domains and Applications ... | 228 |
| 6.10.1 Introduction | 228 |
| 6.11 Proof of Theorem 6.10.1 | 229 |
| 6.12 Application of the Complex Monge–Ampère Equation | 233 |
| 6.13 An Example of David Barrett | 235 |
| 6.14 The Bergman Kernel as a Hilbert Integral | 245 |
| Exercises | 249 |
| 7 Curvature of the Bergman Metric | 251 |
| 7.1 What is the Scaling Method? | 251 |
| 7.2 Higher Dimensional Scaling | 252 |
| 7.2.1 Nonisotropic Scaling | 252 |
| 7.2.2 Normal Convergence of Sets | 254 |
| 7.2.3 Localization | 255 |
| 7.3 Klembeck’s Theorem with C^2 -Stability | 261 |
| 7.3.1 The Main Goal | 261 |
| 7.3.2 The Bergman Metric near Strictly Pseudoconvex Boundary Points | 262 |
| Exercises | 263 |
| 8 Concluding Remarks | 273 |
| Table of Notation | 275 |
| Bibliography | 277 |
| Index | 287 |

Chapter 1

Introductory Ideas

In the early days of functional analysis—the early twentieth century—people did not yet know what a Banach space was nor a Hilbert space. They frequently studied a *particular* complete, infinite-dimensional space from a more abstract point of view. The most common space to be studied in this regard was of course L^2 . It was when Stefan Bergman took a course from Erhard Schmidt on L^2 of the unit interval I that he conceived of the idea of the Bergman space of square-integrable holomorphic functions on the unit disc D . And the rest is history.

It is important for the Bergman theory that his space of holomorphic functions has an inner product structure and that it is complete. The first of these properties follows from the fact that it is a subspace of L^2 ; the second follows from a fundamental inequality that we shall consider in the next section.

1.1 The Bergman Kernel

It is difficult to create an explicit integral formula, with holomorphic reproducing kernel, for holomorphic functions on an arbitrary domain in \mathbb{C}^n .¹ Classical studies which perform such constructions tend to concentrate on domains having a great deal of symmetry (see, for instance, [HUA]). We now examine one of several non-constructive approaches to this problem. This circle of ideas, due to Bergman [BER1] and to Szegő [SZE] (some of the ideas presented here were anticipated by the thesis of Bochner [BOC1]), will later be seen to have profound applications to the boundary regularity of holomorphic mappings.

Bungart [BUN] and Gleason [GLE] have shown that *any* bounded domain in \mathbb{C}^n will have a reproducing kernel for holomorphic functions such that the kernel itself is holomorphic in the free variable. In other words, the formula has the form

¹Here a *domain* is a connected, open set.

$$f(z) = \int_{\Omega} f(\xi)K(z, \xi) dV(\xi),$$

and K is holomorphic in the z variable. Of course Bungart's and Gleason's proofs are highly nonconstructive, and one can say almost nothing about the actual form of the kernel K . The venerable Bochner–Martinelli kernel is easily constructed on any bounded domain with reasonable boundary (just as an application of Stokes's theorem) and the kernel is *explicit*—just like the Cauchy kernel in one complex variable. Also the kernel is the same for every domain. But the Bochner–Martinelli kernel is definitely *not* holomorphic in the free variable. On the other hand, Henkin [HEN], Kerzman [KER1], E. Ramirez [RAMI], and Grauert–Lieb [GRL] have given very explicit constructions of reproducing kernels on strictly pseudoconvex domains (see the definition below). And their kernels *are* holomorphic in the z variable. This matter is treated in [KRA1, Chap. 10].

In fact this last described result was considered to be quite a dramatic advance. For Henkin, Kerzman, Ramirez, and Grauert–Lieb provided us with a fairly explicit kernel, with an explicit and measurable singularity, that can not only reproduce but also create holomorphic functions. Such a kernel is very much like the Cauchy kernel in one complex variable. Thus at least on strictly pseudoconvex domains, we can perform many of the activities to which we are accustomed from the function theory of one complex variable. We can get formulas for derivatives of holomorphic functions, we can analyze power series, we can consider an analogue of the Cauchy transform, and (perhaps most importantly) we can write down solution operators for the $\bar{\partial}$ problem. People were optimistic that these new integral formulas would give a shot in the arm to the theory of function algebras—that they would now be able to study $H^\infty(\Omega)$ and $A(\Omega)$ on a variety of domains in \mathbb{C}^n (see [GAM, Chap. II, IV] for the role model in \mathbb{C}^1). But this turned out to be too difficult.

The Bergman kernel is a canonical kernel that can be defined on any bounded domain. It has wonderful invariance properties and is a powerful tool for geometry and analysis. But it is difficult to calculate explicitly.

In this section we will see some of the invariance properties of the Bergman kernel. This will lead in later sections to the definition of the Bergman metric (in which all biholomorphic mappings become isometries) and to such other canonical constructions as representative coordinates. The Bergman kernel has certain extremal properties that make it a powerful tool in the theory of partial differential equations (see Bergman and Schiffer [BES]). Also the form of the singularity of the Bergman kernel (calculable for some interesting classes of domains) explains many phenomena of the function theory of several complex variables.

Let $\Omega \subseteq \mathbb{C}^n$ be a bounded domain (it is possible, but often tricky, to treat unbounded domains as well). Here a *domain* is a connected, open set. If the domain is smoothly bounded, then we may think of it as specified by a *defining function*:

$$\Omega = \{z \in \mathbb{C}^n : \rho(z) < 0\}.$$

It is customary to require that $\nabla\rho \neq 0$ on $\partial\Omega$. One can demonstrate the existence of a defining function by using the implicit function theorem. See [KRPA2] for the latter and [KRA1] for a detailed consideration of defining functions.

Given a domain Ω as described in the last paragraph and a point $P \in \partial\Omega$, we say that w is a *complex tangent vector* at P and write $w \in \mathcal{T}_P(\partial\Omega)$ if

$$\sum_{j=1}^n \frac{\partial\rho}{\partial z_j}(P)w_j = 0.$$

The point P is said to be *strongly pseudoconvex* if

$$\sum_{j,k=1}^n \frac{\partial^2\rho}{\partial z_j \partial \bar{z}_k}(P)w_j \bar{w}_k > 0$$

for $0 \neq w \in \mathcal{T}_P(\partial\Omega)$. In fact a little elementary analysis shows that we can write the defining property of strong pseudoconvexity as

$$\sum_{j,k=1}^n \frac{\partial^2\rho}{\partial z_j \partial \bar{z}_k}(P)w_j \bar{w}_k \geq C|w|^2$$

and make the estimate uniform when P ranges over a compact, strongly pseudoconvex boundary neighborhood of Ω . Again, the book [KRA1, Chap. 3] has extensive discussion of the notion of strong pseudoconvexity.

Now let us return to the Bergman theory. Let dV denote the Lebesgue volume measure on Ω . Define the *Bergman space*

$$A^2(\Omega) = \left\{ f \text{ holomorphic on } \Omega : \int_{\Omega} |f(z)|^2 dV(z)^{1/2} \equiv \|f\|_{A^2(\Omega)} < \infty \right\}.$$

Of course we equip the Bergman space with the inner product

$$\langle f, g \rangle = \int_{\Omega} f(z)\bar{g}(z) dV(z).$$

Lemma 1.1.1. *Let $K \subseteq \Omega \subseteq \mathbb{C}^n$ be compact. There is a constant $C_K > 0$, depending on K and on n , such that*

$$\sup_{z \in K} |f(z)| \leq C_K \|f\|_{A^2(\Omega)}, \quad \text{all } f \in A^2(\Omega).$$

Proof. Since K is compact, there is an $r(K) = r > 0$ so that, for any $z \in K$, $B(z, r) \subseteq \Omega$. Here $B(z, r)$ is the usual Euclidean ball with center z and radius r .

Therefore for each $z \in K$ and $f \in A^2(\Omega)$, the mean-value property for holomorphic functions implies that

$$\begin{aligned}
|f(z)| &= \left| \frac{1}{V(B(z,r))} \int_{B(z,r)} f(t) dV(t) \right| \\
&= \left| \frac{1}{V(B(z,r))} \int f(t) \chi_{B(z,r)}(t) dV(t) \right| \\
&\leq (V(B(z,r)))^{-1/2} \|f\|_{L^2(B(z,r))} \\
&\leq C(n)r^{-n} \|f\|_{A^2(\Omega)} \\
&\equiv C_K \|f\|_{A^2(\Omega)}. \quad \square
\end{aligned}$$

Lemma 1.1.2. *The space $A^2(\Omega)$ is a Hilbert space with the inner product $\langle f, g \rangle \equiv \int_{\Omega} f(z) \overline{g(z)} dV(z)$.*

Proof. Everything is clear except for completeness. Let $\{f_j\} \subseteq A^2$ be a sequence that is Cauchy in norm. Since L^2 is complete there is an L^2 limit function f . We need to see that f is holomorphic. But Lemma 1.1.1 yields that norm convergence implies normal convergence (i.e., uniform convergence on compact sets). Certainly holomorphic functions are closed under normal limits (just use the Cauchy theory of one complex variable). Therefore f is holomorphic and $A^2(\Omega)$ is complete. \square

Lemma 1.1.3. *For each fixed $z \in \Omega$, the functional*

$$\Phi_z : f \mapsto f(z), \quad f \in A^2(\Omega)$$

is a continuous linear functional on $A^2(\Omega)$.

Proof. This is immediate from Lemma 1.1.1 if we take K to be the singleton $\{z\}$. \square

We may now apply the Riesz representation theorem to see that there is an element $K_z \in A^2(\Omega)$ such that the linear functional Φ_z is represented by inner product with K_z : if $f \in A^2(\Omega)$, then, for all $z \in \Omega$, we have

$$f(z) = \Phi_z(f) = \langle f, K_z \rangle.$$

Definition 1.1.4. *The Bergman kernel is the function $K(z, \zeta) = K_{\Omega}(z, \zeta) \equiv \overline{K_z(\zeta)}$, $z, \zeta \in \Omega$. It has the reproducing property*

$$f(z) = \int_{\Omega} K(z, \zeta) f(\zeta) dV(\zeta), \quad \forall f \in A^2(\Omega).$$

Proposition 1.1.5. *The Bergman kernel $K(z, \zeta)$ is conjugate symmetric: $K(z, \zeta) = \overline{K(\zeta, z)}$.*

Proof. By its very definition, $\overline{K(\zeta, \cdot)} \in A^2(\Omega)$ for each fixed ζ . Therefore the reproducing property of the Bergman kernel gives

$$\int_{\Omega} K(z, t) \overline{K(\zeta, t)} dV(t) = \overline{K(\zeta, z)}.$$

On the other hand,

$$\begin{aligned} \int_{\Omega} K(z, t) \overline{K(\zeta, t)} dV(t) &= \overline{\int_{\Omega} K(\zeta, t) \overline{K(z, t)} dV(t)} \\ &= \overline{\overline{K(z, \zeta)}} = K(z, \zeta). \end{aligned} \quad \square$$

Proposition 1.1.6. *The Bergman kernel is uniquely determined by the properties that it is an element of $A^2(\Omega)$ in z , is conjugate symmetric, and reproduces $A^2(\Omega)$.*

Proof. Let $K'(z, \zeta)$ be another such kernel. Then

$$\begin{aligned} K(z, \zeta) &= \overline{K(\zeta, z)} = \int K'(z, t) \overline{K(\zeta, t)} dV(t) \\ &= \overline{\int K(\zeta, t) \overline{K'(z, t)} dV(t)} \\ &= \overline{\overline{K'(z, \zeta)}} = K'(z, \zeta). \end{aligned} \quad \square$$

Since $L^2(\Omega)$ is a separable Hilbert space then so is its subspace $A^2(\Omega)$. Thus there is a countable, complete orthonormal basis $\{\phi_j\}_{j=1}^{\infty}$ for $A^2(\Omega)$.

Proposition 1.1.7. *Let L be a compact subset of Ω . Then the series*

$$\sum_{j=1}^{\infty} \phi_j(z) \overline{\phi_j(\zeta)}$$

sums uniformly on $L \times L$ to the Bergman kernel $K(z, \zeta)$.

Proof. By the Riesz–Fischer and Riesz representation theorems, we obtain

$$\begin{aligned} \sup_{z \in L} \left(\sum_{j=1}^{\infty} |\phi_j(z)|^2 \right)^{1/2} &= \sup_{z \in L} \left\| \{\phi_j(z)\}_{j=1}^{\infty} \right\|_{\ell^2} \\ &= \sup_{\substack{\|a_j\|_{\ell^2}=1 \\ z \in L}} \left| \sum_{j=1}^{\infty} a_j \phi_j(z) \right| \\ &= \sup_{\substack{\|f\|_{A^2}=1 \\ z \in L}} |f(z)| \\ &\leq C_L. \end{aligned} \quad (1.1.7.1)$$

In the last inequality we have used Lemma 1.1.1. Therefore

$$\sum_{j=1}^{\infty} |\phi_j(z)\overline{\phi_j(\zeta)}| \leq \left(\sum_{j=1}^{\infty} |\phi_j(z)|^2 \right)^{1/2} \left(\sum_{j=1}^{\infty} |\phi_j(\zeta)|^2 \right)^{1/2}$$

and the convergence is uniform over $z, \zeta \in L$. For fixed $z \in \Omega$, (1.1.7.1) shows that $\{\phi_j(z)\}_{j=1}^{\infty} \in \ell^2$. Hence we have that $\sum \phi_j(z)\overline{\phi_j(\zeta)} \in A^2(\Omega)$ as a function of ζ . Let the sum of the series be denoted by $K'(z, \zeta)$. Notice that K' is conjugate symmetric by its very definition. Also, for $f \in A^2(\Omega)$, we have

$$\int K'(\cdot, \zeta) f(\zeta) dV(\zeta) = \sum \hat{f}(j) \phi_j(\cdot) = f(\cdot),$$

where convergence is in the Hilbert space topology. (Here $\hat{f}(j)$ is the j th Fourier coefficient of f with respect to the basis $\{\phi_j\}$.) But Hilbert space convergence dominates pointwise convergence (Lemma 1.1.1) so

$$f(z) = \int K'(z, \zeta) f(\zeta) dV(\zeta), \quad \text{all } f \in A^2(\Omega).$$

Therefore K' is the Bergman kernel. \square

Remark 1.1.8. It is worth noting explicitly that the proof of Proposition 1.1.7 shows that

$$\sum \phi_j(z)\overline{\phi_j(\zeta)}$$

equals the Bergman kernel $K(z, \zeta)$ no matter what the choice of complete orthonormal basis $\{\phi_j\}$ for $A^2(\Omega)$. This can be very useful information in practice. \square

Proposition 1.1.9. *If Ω is a bounded domain in \mathbb{C}^n , then the mapping*

$$P : f \mapsto \int_{\Omega} K(\cdot, \zeta) f(\zeta) dV(\zeta)$$

is the Hilbert space orthogonal projection of $L^2(\Omega, dV)$ onto $A^2(\Omega)$. We call P the Bergman projection.

Proof. Notice that P is idempotent and self-adjoint and that $A^2(\Omega)$ is precisely the set of elements of L^2 that are fixed by P . \square

Definition 1.1.10. Let $\Omega \subseteq \mathbb{C}^n$ be a domain and let $f : \Omega \rightarrow \mathbb{C}^n$ be a *holomorphic mapping*, that is, $f(z) = (f_1(z), \dots, f_n(z))$ with f_1, \dots, f_n holomorphic on Ω . Let $w_j = f_j(z), j = 1, \dots, n$. Then the *holomorphic Jacobian matrix* of f is the matrix

$$J_{\mathbb{C}}f = \frac{\partial(w_1, \dots, w_n)}{\partial(z_1, \dots, z_n)}.$$

Write $z_j = x_j + iy_j, w_k = \xi_k + i\eta_k, j, k = 1, \dots, n$. Then the *real Jacobian matrix* of f is the matrix

$$J_{\mathbb{R}}f = \frac{\partial(\xi_1, \eta_1, \dots, \xi_n, \eta_n)}{\partial(x_1, y_1, \dots, x_n, y_n)}.$$

Proposition 1.1.11. *With notation as in the definition, we have*

$$\det J_{\mathbb{R}}f = |\det J_{\mathbb{C}}f|^2$$

whenever f is a holomorphic mapping.

Proof. We exploit the functoriality of the Jacobian. Let $w = (w_1, \dots, w_n) = f(z) = (f_1(z), \dots, f_n(z))$. Write $z_j = x_j + iy_j, w_j = \xi_j + i\eta_j, j = 1, \dots, n$. Then, using the fact that f is holomorphic,

$$d\xi_1 \wedge d\eta_1 \wedge \dots \wedge d\xi_n \wedge d\eta_n = (\det J_{\mathbb{R}}f(x, y)) dx_1 \wedge dy_1 \wedge \dots \wedge dx_n \wedge dy_n. \quad (1.1.11.1)$$

On the other hand,

$$\begin{aligned} & d\xi_1 \wedge d\eta_1 \wedge \dots \wedge d\xi_n \wedge d\eta_n \\ &= \frac{1}{(2i)^n} d\bar{w}_1 \wedge dw_1 \wedge \dots \wedge d\bar{w}_n \wedge dw_n \\ &= \frac{1}{(2i)^n} \overline{(\det J_{\mathbb{C}}f(z))} (\det J_{\mathbb{C}}f(z)) d\bar{z}_1 \wedge dz_1 \wedge \dots \wedge d\bar{z}_n \wedge dz_n \\ &= |\det J_{\mathbb{C}}f(z)|^2 dx_1 \wedge dy_1 \wedge \dots \wedge dx_n \wedge dy_n. \end{aligned} \quad (1.1.11.2)$$

Equating (1.1.11.1) and (1.1.11.2) gives the result. \square

Exercise for the Reader: Prove Proposition 1.1.11 using only matrix theory (no differential forms). This will give rise to a great appreciation for the theory of differential forms (see Bers [BERS, Chap. 7] for help).

Now we can prove the holomorphic implicit function theorem:

Theorem 1.1.12. Let $f_j(w, z)$, $j = 1, \dots, m$ be holomorphic functions of $(w, z) = ((w_1, \dots, w_m), (z_1, \dots, z_n))$ near a point $(w^0, z^0) \in \mathbb{C}^m \times \mathbb{C}^n$. Assume that

$$f_j(w^0, z^0) = 0, \quad j = 1, \dots, m,$$

and that

$$\det \left(\frac{\partial f_j}{\partial w_k} \right)_{j,k=1}^m \neq 0 \text{ at } (w^0, z^0).$$

Then the system of equations

$$f_j(w, z) = 0, \quad j = 1, \dots, m,$$

has a unique holomorphic solution $w(z)$ in a neighborhood of z^0 that satisfies $w(z^0) = w^0$.

Proof. We rewrite the system of equations as

$$\operatorname{Re} f_j(w, z) = 0, \quad \operatorname{Im} f_j(w, z) = 0$$

for the $2m$ real variables $\operatorname{Re} w_k, \operatorname{Im} w_k, k = 1, \dots, m$. By Proposition 1.1.11, the determinant of the Jacobian over \mathbb{R} of this new system is the modulus squared of the determinant of the Jacobian over \mathbb{C} of the old system. By our hypothesis, this number is nonvanishing at the point (w^0, z^0) . Therefore the classical implicit function theorem (see Rudin [RUD1] or [KRPA2]) implies that there exist C^1 functions $w_k(z), k = 1, \dots, m$, with $w(z^0) = w^0$ and that solve the system. Our job is to show that these functions are in fact holomorphic. When properly viewed, this is purely a problem of geometric algebra:

Applying exterior differentiation to the equations

$$0 = f_j(w(z), z), \quad j = 1, \dots, m,$$

yields that

$$0 = df_j = \sum_{k=1}^m \frac{\partial f_j}{\partial w_k} dw_k + \sum_{k=1}^m \frac{\partial f_j}{\partial z_k} dz_k.$$

There are no $d\bar{z}_j$'s and no $d\bar{w}_k$'s because the f_j 's are holomorphic.

The result now follows from linear algebra only: The hypothesis on the determinant of the matrix $(\partial f_j / \partial w_k)$ implies that we can solve for dw_k in terms of dz_j . Therefore w is a holomorphic function of z . \square

A holomorphic mapping $f : \Omega_1 \rightarrow \Omega_2$ of domains $\Omega_1 \subseteq \mathbb{C}^n, \Omega_2 \subseteq \mathbb{C}^n$ is said to be *biholomorphic* if it is one-to-one, onto, and $\det J_{\mathbb{C}} f(z) \neq 0$ for every $z \in \Omega_1$.

Exercise for the Reader: Use Theorem 1.1.12 to prove that a biholomorphic mapping has a holomorphic inverse (hence the name).

Remark 1.1.13. It is true, but not at all obvious, that the nonvanishing of the Jacobian determinant is a superfluous condition in the definition of “biholomorphic mapping”; that is, the nonvanishing of the Jacobian follows from the univalence of the mapping. A proof of this assertion is sketched in Exercise 37 at the end of [KRA1, Chap. 11]. \square

In what follows we shall frequently denote the Bergman kernel for a given domain Ω by K_{Ω} .

Proposition 1.1.14. *Let Ω_1, Ω_2 be domains in \mathbb{C}^n . Let $f : \Omega_1 \rightarrow \Omega_2$ be biholomorphic. Then*

$$\det J_{\mathbb{C}} f(z) K_{\Omega_2}(f(z), f(\zeta)) \overline{\det J_{\mathbb{C}} f(\zeta)} = K_{\Omega_1}(z, \zeta).$$

Proof. Let $\phi \in A^2(\Omega_1)$. Then, by change of variable,

$$\begin{aligned} & \int_{\Omega_1} \det J_{\mathbb{C}} f(z) K_{\Omega_2}(f(z), f(\zeta)) \overline{\det J_{\mathbb{C}} f(\zeta)} \phi(\zeta) dV(\zeta) \\ &= \int_{\Omega_2} \det J_{\mathbb{C}} f(z) K_{\Omega_2}(f(z), \tilde{\zeta}) \overline{\det J_{\mathbb{C}} f(f^{-1}(\tilde{\zeta}))} \phi(f^{-1}(\tilde{\zeta})) \\ & \quad \times \det J_{\mathbb{R}} f^{-1}(\tilde{\zeta}) dV(\tilde{\zeta}). \end{aligned}$$

By Proposition 1.1.11 this simplifies to

$$\det J_{\mathbb{C}} f(z) \int_{\Omega_2} K_{\Omega_2}(f(z), \tilde{\zeta}) \left\{ \left(\det J_{\mathbb{C}} f(f^{-1}(\tilde{\zeta})) \right)^{-1} \phi \left(f^{-1}(\tilde{\zeta}) \right) \right\} dV(\tilde{\zeta}).$$

By change of variables, the expression in braces $\{ \}$ is an element of $A^2(\Omega_2)$. So the reproducing property of K_{Ω_2} applies and the last line equals

$$= \det J_{\mathbb{C}} f(z) (\det J_{\mathbb{C}} f(z))^{-1} \phi \left(f^{-1}(f(z)) \right) = \phi(z).$$

By the uniqueness of the Bergman kernel, the proposition follows. \square

Proposition 1.1.15. For $z \in \Omega \subset \subset \mathbb{C}^n$ it holds that $K_\Omega(z, z) > 0$.

Proof. Now

$$K_\Omega(z, z) = \sum_{j=1}^{\infty} |\phi_j(z)|^2 \geq 0.$$

If in fact $K(z, z) = 0$ for some z , then $\phi_j(z) = 0$ for all j ; hence, $f(z) = 0$ for every $f \in A^2(\Omega)$. This is absurd. \square

Definition 1.1.16. For any bounded domain $\Omega \subseteq \mathbb{C}^n$, we define a Hermitian metric on Ω by

$$g_{ij}(z) = \frac{\partial^2}{\partial z_i \partial \bar{z}_j} \log K(z, z), \quad z \in \Omega.$$

This means that the square of the length of a tangent vector $\xi = (\xi_1, \dots, \xi_n)$ at a point $z \in \Omega$ is given by

$$|\xi|_{B,z}^2 = \sum_{i,j} g_{ij}(z) \xi_i \bar{\xi}_j.$$

The metric that we have defined is called the *Bergman metric*.

In a Hermitian metric $\{g_{ij}\}$, the length of a C^1 curve $\gamma : [0, 1] \rightarrow \Omega$ is given by

$$\ell(\gamma) = \int_0^1 \left(\sum_{i,j} g_{i,j}(\gamma(t)) \gamma'_i(t) \overline{\gamma'_j(t)} \right)^{1/2} dt.$$

If P, Q are points of Ω , then their distance $d_\Omega(P, Q)$ in the metric is defined to be the infimum of the lengths of all piecewise C^1 curves connecting the two points.

Remark 1.1.17. It is not a priori obvious that the Bergman metric for a bounded domain Ω is given by a positive definite matrix at each point. We now outline a proof of this fact.

First we generate an orthonormal basis for the Bergman space. Fix $z_0 \in \Omega$. Let ϕ_0 be the (unique!) element of A^2 with $\phi_0(z_0)$ real, $\|\phi_0\| = 1$, and $\phi_0(z_0)$ maximal. (Why does such a ϕ_0 exist?) Let ϕ_1 be the (unique) element of A^2 with $\phi_1(z_0) = 0$, $(\partial\phi_1/\partial z_1)(z_0)$ real, $\|\phi_1\| = 1$, and $(\partial\phi_1/\partial z_1)(z_0)$ maximal. (Why does such a ϕ_1 exist?) Now ϕ_1 is orthogonal to ϕ_0 , else ϕ_1 has nonzero projection on ϕ_0 , leading to a contradiction. Continue this process to create an orthogonal system on Ω . Use Taylor series to see that it is complete. This circle of ideas comes from the elegant paper Kobayashi [KOB1].

Now let $\Omega \subseteq \mathbb{C}^n$ be a bounded domain and let (g_{ij}) be its Bergman metric. Use the ideas in the last paragraph to prove that the matrix $(g_{ij}(z))$ is positive definite, each $z \in \Omega$. [*Hint:* The crucial fact is that, for each $z \in \Omega$ and each j , there is an element $f \in A^2(\Omega)$ such that $\partial f / \partial z_j(z) \neq 0$.] \square

- [read The Psychology of Kundalini Yoga: Notes of the Seminar Given in 1932.pdf, azw \(kindle\), epub, doc, mobi](#)
- [Metal Fabrication Technology for Agriculture \(2nd Edition\).pdf, azw \(kindle\), epub, doc, mobi](#)
- [Lost Paradise.pdf, azw \(kindle\), epub, doc, mobi](#)
- [Baby Blueprint Crochet: Irresistible Projects for Little Ones online](#)
- [download Wind Chime Point book](#)

- <http://xn--d1aboelcb1f.xn--p1ai/lib/Fear-Itself--A-Novel.pdf>
- <http://monkeybubblemedia.com/lib/Metal-Fabrication-Technology-for-Agriculture--2nd-Edition-.pdf>
- <http://pittiger.com/lib/Lost-Paradise.pdf>
- <http://www.1973vision.com/?library/Baby-Blueprint-Crochet--Irresistible-Projects-for-Little-Ones.pdf>
- <http://wind-in-herleshausen.de/?freebooks/Wind-Chime-Point.pdf>