


SCHAUM'S
outlines

Calculus

Sixth Edition

1,105 fully solved problems

30 problem-solving videos online 

USE WITH THESE COURSES

Calculus I • Calculus II • Calculus III • AP Calculus • Precalculus



Frank Ayres, Jr., PhD • Elliott Mendelson, PhD



This page intentionally left blank



Visit
Schaums.com to view
Schaum's problem-solving videos
—FREE!


At **Schaums.com**, you'll find videos for
all the most popular Schaum's subjects

Watch and hear instructors explain
problems step by step

Learn valuable problem-solving
techniques

Find out how to tackle common
problem types

Get the benefits of a real
classroom experience



Video Subjects Include: Accounting | Business and Economics | Engineering | Foreign Language | Mathematics | Science

Check out the full range of Schaum resources available from McGraw-Hill Education.



Learn More. Do More.
MHPROFESSIONAL.COM

Preface

The purpose of this book is to help students understand and use the calculus. Everything has been aimed toward making this easier, especially for students with limited background in mathematics or for readers who have forgotten their earlier training in mathematics. The topics covered include all the material of standard courses in elementary and intermediate calculus. The direct and concise exposition typical of the Schaum Outline series has been amplified by a large number of examples, followed by many carefully solved problems. In choosing these problems, we have attempted to anticipate the difficulties that normally beset the beginner. In addition, each chapter concludes with a collection of supplementary exercises with answers.

This sixth edition has enlarged the number of solved problems and supplementary exercises. Moreover, we have made a great effort to go over ticklish points of algebra or geometry that are likely to confuse the student. The author believes that most of the mistakes that students make in a calculus course are not due to a deficient comprehension of the principles of calculus, but rather to their weakness in high-school algebra or geometry. Students are urged to continue the study of each chapter until they are confident about their mastery of the material. A good test of that accomplishment would be their ability to answer the supplementary problems.

The author would like to thank many people who have written to me with corrections and suggestions, in particular Danielle Cinq-Mars, Lawrence Collins, L.D. De Jonge, Konrad Duch, Stephanie Happ, Lindsey Oh, and Stephen B. Soffer. He is also grateful to his editor, Charles Wall, for all his patient help and guidance.

ELLIOTT MENDELSON

This page intentionally left blank

Contents

CHAPTER 1	Linear Coordinate Systems. Absolute Value. Inequalities	1
	Linear Coordinate System Finite Intervals Infinite Intervals Inequalities	
CHAPTER 2	Rectangular Coordinate Systems	9
	Coordinate Axes Coordinates Quadrants The Distance Formula The Midpoint Formulas Proofs of Geometric Theorems	
CHAPTER 3	Lines	18
	The Steepness of a Line The Sign of the Slope Slope and Steepness Equations of Lines A Point–Slope Equation Slope–Intercept Equation Parallel Lines Perpendicular Lines	
CHAPTER 4	Circles	29
	Equations of Circles The Standard Equation of a Circle	
CHAPTER 5	Equations and Their Graphs	37
	The Graph of an Equation Parabolas Ellipses Hyperbolas Conic Sections	
CHAPTER 6	Functions	49
CHAPTER 7	Limits	56
	Limit of a Function Right and Left Limits Theorems on Limits Infinity	
CHAPTER 8	Continuity	66
	Continuous Function	
CHAPTER 9	The Derivative	73
	Delta Notation The Derivative Notation for Derivatives Differentiability	
CHAPTER 10	Rules for Differentiating Functions	79
	Differentiation Composite Functions. The Chain Rule Alternative Formulation of the Chain Rule Inverse Functions Higher Derivatives	

CHAPTER 11 Implicit Differentiation	90
Implicit Functions Derivatives of Higher Order	
CHAPTER 12 Tangent and Normal Lines	93
The Angles of Intersection	
CHAPTER 13 Law of the Mean. Increasing and Decreasing Functions	98
Relative Maximum and Minimum Increasing and Decreasing Functions	
CHAPTER 14 Maximum and Minimum Values	105
Critical Numbers Second Derivative Test for Relative Extrema First Derivative Test Absolute Maximum and Minimum Tabular Method for Finding the Absolute Maximum and Minimum	
CHAPTER 15 Curve Sketching. Concavity. Symmetry	119
Concavity Points of Inflection Vertical Asymptotes Horizontal Asymptotes Symmetry Inverse Functions and Symmetry Even and Odd Functions Hints for Sketching the Graph of $y = f(x)$	
CHAPTER 16 Review of Trigonometry	130
Angle Measure Directed Angles Sine and Cosine Functions	
CHAPTER 17 Differentiation of Trigonometric Functions	139
Continuity of $\cos x$ and $\sin x$ Graph of $\sin x$ Graph of $\cos x$ Other Trigonometric Functions Derivatives Other Relationships Graph of $y = \tan x$ Graph of $y = \sec x$ Angles Between Curves	
CHAPTER 18 Inverse Trigonometric Functions	152
The Derivative of $\sin^{-1}x$ The Inverse Cosine Function The Inverse Tangent Function	
CHAPTER 19 Rectilinear and Circular Motion	161
Rectilinear Motion Motion Under the Influence of Gravity Circular Motion	
CHAPTER 20 Related Rates	167
CHAPTER 21 Differentials. Newton's Method	173
The Differential Newton's Method	
CHAPTER 22 Antiderivatives	181
Laws for Antiderivatives	

CHAPTER 23 The Definite Integral. Area Under a Curve	190
Sigma Notation Area Under a Curve Properties of the Definite Integral	
CHAPTER 24 The Fundamental Theorem of Calculus	198
Mean-Value Theorem for Integrals Average Value of a Function on a Closed Interval Fundamental Theorem of Calculus Change of Variable in a Definite Integral	
CHAPTER 25 The Natural Logarithm	206
The Natural Logarithm Properties of the Natural Logarithm	
CHAPTER 26 Exponential and Logarithmic Functions	214
Properties of e^x The General Exponential Function General Logarithmic Functions	
CHAPTER 27 L'Hôpital's Rule	222
L'Hôpital's Rule Indeterminate Type $0 \cdot \infty$ Indeterminate Type $\infty - \infty$ Indeterminate Types 0^0 , ∞^0 , and 1^∞	
CHAPTER 28 Exponential Growth and Decay	230
Half-Life	
CHAPTER 29 Applications of Integration I: Area and Arc Length	235
Area Between a Curve and the y Axis Areas Between Curves Arc Length	
CHAPTER 30 Applications of Integration II: Volume	244
Disk Formula Washer Method Cylindrical Shell Method Difference of Shells Formula Cross-Section Formula (Slicing Formula)	
CHAPTER 31 Techniques of Integration I: Integration by Parts	259
CHAPTER 32 Techniques of Integration II: Trigonometric Integrands and Trigonometric Substitutions	266
Trigonometric Integrands Trigonometric Substitutions	
CHAPTER 33 Techniques of Integration III: Integration by Partial Fractions	279
Method of Partial Fractions	
CHAPTER 34 Techniques of Integration IV: Miscellaneous Substitutions	288

CHAPTER 35 Improper Integrals	293
Infinite Limits of Integration Discontinuities of the Integrand	
CHAPTER 36 Applications of Integration III: Area of a Surface of Revolution	301
CHAPTER 37 Parametric Representation of Curves	307
Parametric Equations Arc Length for a Parametric Curve	
CHAPTER 38 Curvature	312
Derivative of Arc Length Curvature The Radius of Curvature The Circle of Curvature The Center of Curvature The Evolute	
CHAPTER 39 Plane Vectors	321
Scalars and Vectors Sum and Difference of Two Vectors Components of a Vector Scalar Product (or Dot Product) Scalar and Vector Projections Differentiation of Vector Functions	
CHAPTER 40 Curvilinear Motion	332
Velocity in Curvilinear Motion Acceleration in Curvilinear Motion Tangential and Normal Components of Acceleration	
CHAPTER 41 Polar Coordinates	339
Polar and Rectangular Coordinates Some Typical Polar Curves Angle of Inclination Points of Intersection Angle of Intersection The Derivative of the Arc Length Curvature	
CHAPTER 42 Infinite Sequences	352
Infinite Sequences Limit of a Sequence Monotonic Sequences	
CHAPTER 43 Infinite Series	360
Geometric Series	
CHAPTER 44 Series with Positive Terms. The Integral Test. Comparison Tests	366
Series of Positive Terms	
CHAPTER 45 Alternating Series. Absolute and Conditional Convergence. The Ratio Test	375
Alternating Series	
CHAPTER 46 Power Series	383
Power Series Uniform Convergence	

CHAPTER 47 Taylor and Maclaurin Series. Taylor's Formula with Remainder	396
Taylor and Maclaurin Series Applications of Taylor's Formula with Remainder	
CHAPTER 48 Partial Derivatives	405
Functions of Several Variables Limits Continuity Partial Derivatives Partial Derivatives of Higher Order	
CHAPTER 49 Total Differential. Differentiability. Chain Rules	414
Total Differential Differentiability Chain Rules Implicit Differentiation	
CHAPTER 50 Space Vectors	426
Vectors in Space Direction Cosines of a Vector Determinants Vector Perpendicular to Two Vectors Vector Product of Two Vectors Triple Sca- lar Product Triple Vector Product The Straight Line The Plane	
CHAPTER 51 Surfaces and Curves in Space	441
Planes Spheres Cylindrical Surfaces Ellipsoid Elliptic Paraboloid Elliptic Cone Hyperbolic Paraboloid Hyperboloid of One Sheet Hyperbo- loid of Two Sheets Tangent Line and Normal Plane to a Space Curve Tangent Plane and Normal Line to a Surface Surface of Revolution	
CHAPTER 52 Directional Derivatives. Maximum and Minimum Values	452
Directional Derivatives Relative Maximum and Minimum Values Absolute Maximum and Minimum Values	
CHAPTER 53 Vector Differentiation and Integration	460
Vector Differentiation Space Curves Surfaces The Operation ∇ Divergence and Curl Integration Line Integrals	
CHAPTER 54 Double and Iterated Integrals	474
The Double Integral The Iterated Integral	
CHAPTER 55 Centroids and Moments of Inertia of Plane Areas	481
Plane Area by Double Integration Centroids Moments of Inertia	
CHAPTER 56 Double Integration Applied to Volume Under a Surface and the Area of a Curved Surface	489
CHAPTER 57 Triple Integrals	498
Cylindrical and Spherical Coordinates The Triple Integral Evaluation of Triple Integrals Centroids and Moments of Inertia	
CHAPTER 58 Masses of Variable Density	510

CHAPTER 59 Differential Equations of First and Second Order	516	
Separable Differential Equations	Homogeneous Functions	Integrating Factors
Second-Order Equations		
Appendix A	527	
Appendix B	528	
Index	529	

Linear Coordinate Systems. Absolute Value. Inequalities

Linear Coordinate System

A linear coordinate system is a graphical representation of the real numbers as the points of a straight line. To each number corresponds one and only one point, and to each point corresponds one and only one number.

To set up a linear coordinate system on a given line: (1) select any point of the line as the *origin* and let that point correspond to the number 0; (2) choose a positive direction on the line and indicate that direction by an arrow; (3) choose a fixed distance as a unit of measure. If x is a positive number, find the point corresponding to x by moving a distance of x units from the origin in the positive direction. If x is negative, find the point corresponding to x by moving a distance of $-x$ units from the origin in the negative direction. (For example, if $x = -2$, then $-x = 2$ and the corresponding point lies 2 units from the origin in the negative direction.) See Fig. 1-1.




Fig. 1-1

The number assigned to a point by a coordinate system is called the *coordinate* of that point. We often will talk as if there is no distinction between a point and its coordinate. Thus, we might refer to “the point 3” rather than to “the point with coordinate 3.”

The absolute value $|x|$ of a number x is defined as follows:

$$|x| = \begin{cases} x & \text{if } x \text{ is zero or a positive number} \\ -x & \text{if } x \text{ is a negative number} \end{cases}$$

For example, $|4| = 4$, $|-3| = -(-3) = 3$, and $|0| = 0$. Notice that, if x is a negative number, then $-x$ is positive. Thus, $|x| \geq 0$ for all x .

The following properties hold for any numbers x and y .

- (1.1) $| -x | = | x |$
 When $x = 0$, $| -x | = | -0 | = | 0 | = | x |$.
 When $x > 0$, $-x < 0$ and $| -x | = -(-x) = x = | x |$.
 When $x < 0$, $-x > 0$, and $| -x | = -x = | x |$.
- (1.2) $| x - y | = | y - x |$
 This follows from (1.1), since $y - x = -(x - y)$.
- (1.3) $| x | = c$ implies that $x = \pm c$.
 For example, if $| x | = 2$, then $x = \pm 2$. For the proof, assume $| x | = c$.
 If $x \geq 0$, $x = | x | = c$. If $x < 0$, $-x = | x | = c$; then $x = -(-x) = -c$.
- (1.4) $| x |^2 = x^2$
 If $x \geq 0$, $| x | = x$ and $| x |^2 = x^2$. If $x \leq 0$, $| x | = -x$ and $| x |^2 = (-x)^2 = x^2$.
- (1.5) $| xy | = | x | \cdot | y |$
 By (1.4), $(| xy |)^2 = (xy)^2 = x^2 y^2 = | x |^2 | y |^2 = (| x | \cdot | y |)^2$. Since absolute values are nonnegative, taking square roots yields $| xy | = | x | \cdot | y |$.

$$(1.6) \quad \left| \frac{x}{y} \right| = \frac{|x|}{|y|} \text{ if } y \neq 0$$

By (1.5), $|y| \left| \frac{x}{y} \right| = \left| y \cdot \frac{x}{y} \right| = |x|$. Divide by $|y|$.

$$(1.7) \quad |x| = |y| \text{ implies that } x = \pm y$$

Assume $|x| = |y|$. If $y = 0$, $|x| = |0| = 0$ and (1.3) yields $x = 0$. If $y \neq 0$, then by (1.6),

$$\left| \frac{x}{y} \right| = \frac{|x|}{|y|} = 1$$

So, by (1.3), $x/y = \pm 1$. Hence, $x = \pm y$.

$$(1.8) \quad \text{Let } c \geq 0. \text{ Then } |x| \leq c \text{ if and only if } -c \leq x \leq c. \text{ See Fig. 1-2.}$$

Assume $x \geq 0$. Then $|x| = x$. Also, since $c \geq 0$, $-c \leq 0 \leq x$. So, $|x| \leq c$ if and only if $-c \leq x \leq c$. Now assume $x < 0$. Then $|x| = -x$. Also, $x < 0 \leq c$. Moreover, $-x \leq c$ if and only if $-c \leq x$. (Multiplying or dividing an equality by a negative number reverses the inequality.) Hence, $|x| \leq c$ if and only if $-c \leq x \leq c$.

$$(1.9) \quad \text{Let } c \geq 0. \text{ Then } |x| < c \text{ if and only if } -c < x < c. \text{ See Fig. 1-2. The reasoning here is similar to that for (1.8).}$$



Fig. 1-2

$$(1.10) \quad -|x| \leq x \leq |x|$$

If $x \geq 0$, $x = |x|$. If $x < 0$, $|x| = -x$ and, therefore, $x = -|x|$.

$$(1.11) \quad |x + y| \leq |x| + |y| \text{ (triangle inequality)}$$

By (1.8), $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$. Adding, we obtain $-(|x| + |y|) \leq x + y \leq |x| + |y|$. Then $|x + y| \leq |x| + |y|$ by (1.8). [In (1.8), replace c by $|x| + |y|$ and x by $x + y$.]

Let a coordinate system be given on a line. Let P_1 and P_2 be points on the line having coordinates x_1 and x_2 . See Fig. 1-3. Then:

$$(1.12) \quad |x_1 - x_2| = P_1 P_2 = \text{distance between } P_1 \text{ and } P_2.$$

This is clear when $0 < x_1 < x_2$ and when $x_1 < x_2 < 0$. When $x_1 < 0 < x_2$, and if we denote the origin by O , then $P_1 P_2 = P_1 O + O P_2 = (-x_1) + x_2 = x_2 - x_1 = |x_2 - x_1| = |x_1 - x_2|$.

As a special case of (1.12), when P_2 is the origin (and $x_2 = 0$):

$$(1.13) \quad |x_1| = \text{distance between } P_1 \text{ and the origin.}$$



Fig. 1-3

Finite Intervals

Let $a < b$.

The *open interval* (a, b) is defined to be the set of all numbers between a and b , that is, the set of all x such that $a < x < b$. We shall use the term *open interval* and the notation (a, b) also for all the points between the points with coordinates a and b on a line. Notice that the open interval (a, b) does not contain the *endpoints* a and b . See Fig. 1-4.

The *closed interval* $[a, b]$ is defined to be the set of all numbers between a and b or equal to a or b , that is, the set of all x such that $a \leq x \leq b$. As in the case of open intervals, we extend the terminology and notation to points. Notice that the closed interval $[a, b]$ contains both endpoints a and b . See Fig. 1-4.



Fig. 1-4

By a *half-open interval* we mean an open interval (a, b) together with one of its endpoints. There are two such intervals: $[a, b)$ is the set of all x such that $a \leq x < b$, and $(a, b]$ is the set of all x such that $a < x \leq b$.

Infinite Intervals

Let (a, ∞) denote the set of all x such that $a < x$.
 Let $[a, \infty)$ denote the set of all x such that $a \leq x$.
 Let $(-\infty, b)$ denote the set of all x such that $x < b$.
 Let $(-\infty, b]$ denote the set of all x such that $x \leq b$.

Inequalities

Any inequality, such as $2x - 3 > 0$ or $5 < 3x + 10 \leq 16$, determines an interval. To solve an inequality means to determine the corresponding interval of numbers that satisfy the inequality.

EXAMPLE 1.1: Solve $2x - 3 > 0$.

$$\begin{aligned} 2x - 3 &> 0 \\ 2x &> 3 && \text{(Adding 3)} \\ x &> \frac{3}{2} && \text{(Dividing by 2)} \end{aligned}$$

Thus, the corresponding interval is $(\frac{3}{2}, \infty)$.

EXAMPLE 1.2: Solve $5 < 3x + 10 \leq 16$.

$$\begin{aligned} 5 &< 3x + 10 \leq 16 \\ -5 &< 3x \leq 6 && \text{(Subtracting 10)} \\ -\frac{5}{3} &< x \leq 2 && \text{(Dividing by 3)} \end{aligned}$$

Thus, the corresponding interval is $(-\frac{5}{3}, 2]$.

EXAMPLE 1.3: Solve $-2x + 3 < 7$.

$$\begin{aligned} -2x + 3 &< 7 \\ -2x &< 4 && \text{(Subtracting 3)} \\ x &> -2 && \text{(Dividing by } -2) \end{aligned}$$

(Recall that dividing by a negative number reverses an inequality.) Thus, the corresponding interval is $(-2, \infty)$.

SOLVED PROBLEMS

- Describe and diagram the following intervals, and write their interval notation, (a) $-3 < x < 5$; (b) $2 \leq x \leq 6$; (c) $-4 < x \leq 0$; (d) $x > 5$; (e) $x \leq 2$; (f) $3x - 4 \leq 8$; (g) $1 < 5 - 3x < 11$.
 (a) All numbers greater than -3 and less than 5 ; the interval notation is $(-3, 5)$:



- (b) All numbers equal to or greater than 2 and less than or equal to 6; $[2, 6]$:



- (c) All numbers greater than -4 and less than or equal to 0 ; $(-4, 0]$:



- (d) All numbers greater than 5 ; $(5, \infty)$:



- (e) All numbers less than or equal to 2 ; $(-\infty, 2]$:



- (f) $3x - 4 \leq 8$ is equivalent to $3x \leq 12$ and, therefore, to $x \leq 4$. Thus, we get $(-\infty, 4]$:



- (g) $1 < 5 - 3x < 11$
 $-4 < -3x < 6$ (Subtracting 5)
 $-2 < x < \frac{4}{3}$ (Dividing by -3 ; note the reversal of inequalities)

Thus, we obtain $(-2, \frac{4}{3})$:




2. Describe and diagram the intervals determined by the following inequalities, (a) $|x| < 2$; (b) $|x| > 3$; (c) $|x - 3| < 1$; (d) $|x - 2| < \delta$ where $\delta > 0$; (e) $|x + 2| \leq 3$; (f) $0 < |x - 4| < \delta$ where $\delta > 0$.

- (a) By property (1.9), this is equivalent to $-2 < x < 2$, defining the open interval $(-2, 2)$.



- (b) By property (1.8), $|x| \leq 3$ is equivalent to $-3 \leq x \leq 3$. Taking negations, $|x| > 3$ is equivalent to $x < -3$ or $x > 3$, which defines the union of the intervals $(-\infty, -3)$ and $(3, \infty)$.



- (c) By property (1.12), this says that the distance between x and 3 is less than 1 , which is equivalent to $2 < x < 4$. This defines the open interval $(2, 4)$.



We can also note that $|x - 3| < 1$ is equivalent to $-1 < x - 3 < 1$. Adding 3 , we obtain $2 < x < 4$.

- (d) This is equivalent to saying that the distance between x and 2 is less than δ , or that $2 - \delta < x < 2 + \delta$, which defines the open interval $(2 - \delta, 2 + \delta)$. This interval is called the δ -neighborhood of 2 :



- (e) $|x + 2| < 3$ is equivalent to $-3 < x + 2 < 3$. Subtracting 2, we obtain $-5 < x < 1$, which defines the open interval $(-5, 1)$:




- (f) The inequality $|x - 4| < \delta$ determines the interval $4 - \delta < x < 4 + \delta$. The additional condition $0 < |x - 4|$ tells us that $x \neq 4$. Thus, we get the union of the two intervals $(4 - \delta, 4)$ and $(4, 4 + \delta)$. The result is called the *deleted δ -neighborhood* of 4:



3. Describe and diagram the intervals determined by the following inequalities, (a) $|5 - x| \leq 3$; (b) $|2x - 3| < 5$; (c) $|1 - 4x| < \frac{1}{2}$.

- (a) Since $|5 - x| = |x - 5|$, we have $|x - 5| \leq 3$, which is equivalent to $-3 \leq x - 5 \leq 3$. Adding 5, we get $2 \leq x \leq 8$, which defines the closed interval $[2, 8]$:



- (b) $|2x - 3| < 5$ is equivalent to $-5 < 2x - 3 < 5$. Adding 3, we have $-2 < 2x < 8$; then dividing by 2 yields $-1 < x < 4$, which defines the open interval $(-1, 4)$:



- (c) Since $|1 - 4x| = |4x - 1|$, we have $|4x - 1| < \frac{1}{2}$, which is equivalent to $-\frac{1}{2} < 4x - 1 < \frac{1}{2}$. Adding 1, we get $\frac{1}{2} < 4x < \frac{3}{2}$. Dividing by 4, we obtain $\frac{1}{8} < x < \frac{3}{8}$, which defines the open interval $(\frac{1}{8}, \frac{3}{8})$:




4. Solve the inequalities: (a) $18x - 3x^2 > 0$; (b) $(x + 3)(x - 2)(x - 4) < 0$; (c) $(x + 1)^2(x - 3) > 0$, and diagram the solutions.

- (a) Set $18x - 3x^2 = 3x(6 - x) = 0$, obtaining $x = 0$ and $x = 6$. We need to determine the sign of $18x - 3x^2$ on each of the intervals $x < 0$, $0 < x < 6$, and $x > 6$, to determine where $18x - 3x^2 > 0$. Note that it is negative when $x < 0$ (since x is negative and $6 - x$ is positive). It becomes positive when we pass from left to right through 0 (since x changes sign but $6 - x$ remains positive), and it becomes negative when we pass through 6 (since x remains positive but $6 - x$ changes to negative). Hence, it is positive when and only when $0 < x < 6$.



- (b) The crucial points are $x = -3$, $x = 2$, and $x = 4$. Note that $(x + 3)(x - 2)(x - 4)$ is negative for $x < -3$ (since each of the factors is negative) and that it changes sign when we pass through each of the crucial points. Hence, it is negative for $x < -3$ and for $2 < x < 4$:



- (c) Note that $(x + 1)$ is always positive (except at $x = -1$, where it is 0). Hence $(x + 1)^2(x - 3) > 0$ when and only when $x - 3 > 0$, that is, for $x > 3$:



5. Solve $|3x - 7| = 8$.

By (1.3), $|3x - 7| = 8$ if and only if $3x - 7 = \pm 8$. Thus, we need to solve $3x - 7 = 8$ and $3x - 7 = -8$. Hence, we get $x = 5$ or $x = -\frac{1}{3}$.

6. Solve $\frac{2x+1}{x+3} > 3$.

Case 1: $x+3 > 0$. Multiply by $x+3$ to obtain $2x+1 > 3x+9$, which reduces to $-8 > x$. However, since $x+3 > 0$, it must be that $x > -3$. Thus, this case yields no solutions.

Case 2: $x+3 < 0$. Multiply by $x+3$ to obtain $2x+1 < 3x+9$. (Note that the inequality is reversed, since we multiplied by a negative number.) This yields $-8 < x$. Since $x+3 < 0$, we have $x < -3$. Thus, the only solutions are $-8 < x < -3$.

7. Solve $\left| \frac{2}{x} - 3 \right| < 5$.

The given inequality is equivalent to $-5 < \frac{2}{x} - 3 < 5$. Add 3 to obtain $-2 < 2/x < 8$, and divide by 2 to get $-1 < 1/x < 4$.

Case 1: $x > 0$. Multiply by x to get $-x < 1 < 4x$. Then $x > \frac{1}{4}$ and $x > -1$; these two inequalities are equivalent to the single inequality $x > \frac{1}{4}$.

Case 2: $x < 0$. Multiply by x to obtain $-x > 1 > 4x$. (Note that the inequalities have been reversed, since we multiplied by the negative number x .) Then $x < \frac{1}{4}$ and $x < -1$. These two inequalities are equivalent to $x < -1$.

Thus, the solutions are $x > \frac{1}{4}$ or $x < -1$, the union of the two infinite intervals $(\frac{1}{4}, \infty)$ and $(-\infty, -1)$.

8. Solve $|2x - 5| \geq 3$.

Let us first solve the negation $|2x - 5| < 3$. The latter is equivalent to $-3 < 2x - 5 < 3$. Add 5 to obtain $2 < 2x < 8$, and divide by 2 to obtain $1 < x < 4$. Since this is the solution of the negation, the original inequality has the solution $x \leq 1$ or $x \geq 4$.

9. Solve: $x^2 < 3x + 10$.

$$\begin{aligned} x^2 &< 3x + 10 \\ x^2 - 3x - 10 &< 0 && \text{(Subtract } 3x + 10\text{)} \\ (x - 5)(x + 2) &< 0 \end{aligned}$$

The crucial numbers are -2 and 5 . $(x - 5)(x + 2) > 0$ when $x < -2$ (since both $x - 5$ and $x + 2$ are negative); it becomes negative as we pass through -2 (since $x + 2$ changes sign); and then it becomes positive as we pass through 5 (since $x - 5$ changes sign). Thus, the solutions are $-2 < x < 5$.

SUPPLEMENTARY PROBLEMS

10. Describe and diagram the set determined by each of the following conditions:

- | | |
|-----------------------------|---------------------------------|
| (a) $-5 < x < 0$ | (b) $x \leq 0$ |
| (c) $-2 \leq x < 3$ | (d) $x \geq 1$ |
| (e) $ x < 3$ | (f) $ x \geq 5$ |
| (g) $ x - 2 < \frac{1}{2}$ | (h) $ x - 3 > 1$ |
| (i) $0 < x - 2 < 1$ | (j) $0 < x + 3 < \frac{1}{4}$ |
| (k) $ x - 2 \geq 1$. | |

Ans. (e) $-3 < x < 3$; (f) $x \geq 5$ or $x \leq -5$; (g) $\frac{3}{2} < x < \frac{5}{2}$; (h) $x > -2$ or $x < -4$; (i) $x \neq 2$ and $1 < x < 3$;
(j) $-\frac{13}{4} < x < -\frac{11}{4}$; (k) $x \geq 3$ or $x \leq 1$

11. Describe and diagram the set determined by each of the following conditions:

- (a) $|3x - 7| < 2$
 (b) $|4x - 11| \geq 1$
 (c) $\left| \frac{x}{3} - 2 \right| \leq 4$

(d) $\left| \frac{3}{x} - 2 \right| \leq 4$

(e) $\left| 2 + \frac{1}{x} \right| > 1$

(f) $\left| \frac{4}{x} \right| < 3$

Ans. (a) $\frac{5}{3} < x < 3$; (b) $x \geq \frac{1}{2}$ or $x \leq 0$; (c) $-6 \leq x \leq 18$; (d) $x \leq -\frac{3}{2}$ or $x \geq \frac{1}{2}$; (e) $x > 0$ or $x < -1$ or $-\frac{1}{3} < x < 0$;
 (f) $x > \frac{4}{3}$ or $x < -\frac{4}{3}$

12. Describe and diagram the set determined by each of the following conditions:

(a) $x(x-5) < 0$

(b) $(x-2)(x-6) > 0$

(c) $(x+1)(x-2) < 0$

(d) $x(x-2)(x+3) > 0$

(e) $(x+2)(x+3)(x+4) < 0$

(f) $(x-1)(x+1)(x-2)(x+3) > 0$

(g) $(x-1)^2(x+4) > 0$

(h) $(x-3)(x+5)(x-4)^2 < 0$

(i) $(x-2)^3 > 0$

(j) $(x+1)^3 < 0$

(k) $(x-2)^3(x+1) < 0$

(l) $(x-1)^3(x+1)^4 < 0$

(m) $(3x-1)(2x+3) > 0$

(n) $(x-4)(2x-3) < 0$

Ans. (a) $0 < x < 5$; (b) $x > 6$ or $x < 2$; (c) $-1 < x < 2$; (d) $x > 2$ or $-3 < x < 0$; (e) $-3 < x < -2$ or $x < -4$;
 (f) $x > 2$ or $-1 < x < 1$ or $x < -3$; (g) $x > -4$ and $x \neq 1$; (h) $-5 < x < 3$; (i) $x > 2$; (j) $x < -1$;
 (k) $-1 < x < 2$; (l) $x < 1$ and $x \neq -1$; (m) $x > \frac{1}{3}$ or $x < -\frac{3}{2}$; (n) $\frac{3}{2} < x < 4$

13. Describe and diagram the set determined by each of the following conditions:

(a) $x^2 < 4$

(b) $x^2 \geq 9$

(c) $(x-2)^2 \leq 16$

(d) $(2x+1)^2 > 1$

(e) $x^2 + 3x - 4 > 0$

(f) $x^2 + 6x + 8 \leq 0$

(g) $x^2 < 5x + 14$

(h) $2x^2 > x + 6$

(i) $6x^2 + 13x < 5$

(j) $x^3 + 3x^2 > 10x$

Ans. (a) $-2 < x < 2$; (b) $x \geq 3$ or $x \leq -3$; (c) $-2 \leq x \leq 6$; (d) $x > 0$ or $x < -1$; (e) $x > 1$ or $x < -4$; (f) $-4 \leq x \leq -2$;
 (g) $-2 < x < 7$; (h) $x > 2$ or $x < -\frac{3}{2}$; (i) $-\frac{5}{2} < x < \frac{1}{3}$; (j) $-5 < x < 0$ or $x > 2$

14. Solve: (a) $-4 < 2 - x < 7$ (b) $\frac{2x-1}{x} < 3$ (c) $\frac{x}{x+2} < 1$
 (d) $\frac{3x-1}{2x+3} > 3$ (e) $\left| \frac{2x-1}{x} \right| > 2$ (f) $\left| \frac{x}{x+2} \right| \leq 2$

Ans. (a) $-5 < x < 6$; (b) $x > 0$ or $x < -1$; (c) $x > -2$; (d) $-\frac{10}{3} < x < \frac{3}{2}$; (e) $x < 0$ or $0 < x < \frac{1}{4}$; (f) $x \leq -4$ or $x \geq -1$

15. Solve:

- (a) $|4x - 5| = 3$
- (b) $|x + 6| = 2$
- (c) $|3x - 4| = |2x + 1|$
- (d) $|x + 1| = |x + 2|$
- (e) $|x + 1| = 3x - 1$
- (f) $|x + 1| < |3x - 1|$
- (g) $|3x - 4| \geq |2x + 1|$

Ans. (a) $x = 2$ or $x = \frac{1}{2}$; (b) $x = -4$ or $x = -8$; (c) $x = 5$ or $x = \frac{5}{3}$; (d) $x = -\frac{3}{2}$; (e) $x = 1$; (f) $x > 1$ or $x < 0$;
(g) $x \geq 5$ or $x \leq \frac{3}{5}$

16. Prove:

- (a) $|x^2| = |x|^2$;
 - (b) $|x^n| = |x|^n$ for every integer n ;
 - (c) $|x| = \sqrt{x^2}$;
 - (d) $|x - y| \leq |x| + |y|$;
 - (e) $|x - y| \geq ||x| - |y||$
- [Hint: In (e), prove that $|x - y| \geq |x| - |y|$ and $|x - y| \geq |y| - |x|$.]

CHAPTER 2

Rectangular Coordinate Systems

Coordinate Axes

In any plane \mathcal{P} , choose a pair of perpendicular lines. Let one of the lines be horizontal. Then the other line must be vertical. The horizontal line is called the x axis, and the vertical line the y axis. (See Fig. 2-1.)




Fig. 2-1

Now choose linear coordinate systems on the x axis and the y axis satisfying the following conditions: The origin for each coordinate system is the point O at which the axes intersect. The x axis is directed from left to right, and the y axis from bottom to top. The part of the x axis with positive coordinates is called the *positive x axis*, and the part of the y axis with positive coordinates is called the *positive y axis*.

We shall establish a correspondence between the points of the plane \mathcal{P} and pairs of real numbers.

Coordinates

Consider any point P of the plane (Fig. 2-1). The vertical line through P intersects the x axis at a unique point; let a be the coordinate of this point on the x axis. The number a is called the x coordinate of P (or the *abscissa* of P). The horizontal line through P intersects the y axis at a unique point; let b be the coordinate of this point on the y axis. The number b is called the y coordinate of P (or the *ordinate* of P). In this way, every point P has a unique pair (a, b) of real numbers associated with it. Conversely, every pair (a, b) of real numbers is associated with a unique point in the plane.

The coordinates of several points are shown in Fig. 2-2. For the sake of simplicity, we have limited them to integers.




Fig. 2-2

EXAMPLE 2.1: In the coordinate system of Fig. 2-3, to find the point having coordinates $(2, 3)$, start at the origin, move two units to the *right*, and then three units *upward*.




Fig. 2-3

To find the point with coordinates $(-4, 2)$, start at the origin, move four units to the *left*, and then two units *upward*.

To find the point with coordinates $(-3, -1)$, start at the origin, move three units to the *left*, and then one unit *downward*.

The order of these moves is not important. Hence, for example, the point $(2, 3)$ can also be reached by starting at the origin, moving three units *upward*, and then two units to the *right*.

Quadrants

Assume that a coordinate system has been established in the plane \mathcal{P} . Then the whole plane \mathcal{P} , with the exception of the coordinate axes, can be divided into four equal parts, called *quadrants*. All points with both coordinates positive form the first quadrant, called quadrant I, in the upper-right-hand corner (see Fig. 2-4).

- **[read Memory of Water](#)**
- [read online Saving Grace \(Serve and Protect, Book 2\) pdf, azw \(kindle\), epub](#)
- [download online Your Credit Score: How to Improve the 3-Digit Number That Shapes Your Financial Future \(4th Edition\) \(Liz Pulliam Weston\) pdf, azw \(kindle\), epub, doc, mobi](#)
- [A Guide to Flexible Dieting pdf, azw \(kindle\), epub, doc, mobi](#)
- [download Taking It Off, Putting It on: Women in the Strip Trade](#)

- <http://monkeybubblemedia.com/lib/Ficciones.pdf>
- <http://www.1973vision.com/?library/Becoming-Bea--The-Courtships-of-Lancaster-County--Book-4-.pdf>
- <http://aseasonedman.com/ebooks/The-Black-Dahlia.pdf>
- <http://cambridgebrass.com/?freebooks/Implementing-Cloud-Storage-with-OpenStack-Swift.pdf>
- <http://nexson.arzamaszev.com/library/Bagel-Greats---Delicious-Bagel-Recipes.pdf>